

THE ACCRETION RATES AND SPECTRAL ENERGY DISTRIBUTIONS OF BL LACERTAE OBJECTS

JIAN-MIN WANG^{1,2,3}, RÜDIGER STAUBERT¹ AND LUIS C. HO⁴*Accepted by the Astrophysical Journal on 12/July, 2002.*

ABSTRACT

We investigate the relationship between accretion rates and the spectral energy distributions (SEDs) of BL Lac objects, using a sample of objects for which published information on the host galaxies, emission-line luminosities, and peak frequencies and luminosities of their SEDs are available. The sample is composed of 43 BL Lac objects which have a relatively continuous distribution of peak frequencies. Under the assumption that the observed emission lines are photoionized by the central accretion disk, we use the line luminosities to estimate the accretion luminosities and hence accretion rates. We find that low frequency-peaked BL Lac objects (LBLs) span a wide range of accretion rates, whereas high frequency-peaked BL Lac objects (HBLs) cover a more restricted range of lower values. There appears to be a continuous distribution of accretion rates between the two subclasses of BL Lac objects. We find that the peak frequency of the SED, ν_{pk} , correlates with the accretion rate, approximately with the form $\nu_{\text{pk}} \propto \Lambda^{-3}$ in HBLs and $\nu_{\text{pk}} \propto \Lambda^{-0.25}$ in LBLs, where $\Lambda \equiv L_{\text{lines}}/c^2$. The peak luminosity of the SED is also correlated with Λ . These results suggest that the accretion rate influences the shape of the SED in BL Lac objects. They also support models which couple the jet and the accretion disk. We present a physical scenario to account for the empirical trends.

Subject headings: accretion, accretion disks — black hole physics — BL Lacertae objects: general — galaxies: active — galaxies: jets — galaxies: nuclei

1. INTRODUCTION

The spectral energy distributions (SEDs) of BL Lac objects can be largely characterized by two broad peaks. The low and high energy peaks are commonly attributed to synchrotron emission and inverse Compton scattering, respectively. Depending on the relative strengths of the two peaks, it is now customary to divide BL Lac objects into low frequency-peaked (LBL) and high frequency-peaked (HBL) sources (Giommi & Padovani 1994). Typically, HBLs are less variable, less polarized, and contain less dominant radio cores than LBLs (e.g., Laurent-Muehleisen et al. 1993; Perlman & Stocke 1993; Januzzi, Smith & Elston 1994). However, the physical origin of the differences between the two classes of BL Lac objects remains a matter of debate.

According to the unification scheme for BL Lac objects, LBLs and HBLs are observed at different viewing angles (see review by Urry & Padovani 1995). On the other hand, Sambruna, Maraschi, & Urry (1996) have shown that the typical multiwaveband SED of an HBL cannot be produced from the spectrum of an LBL simply by changing the viewing angle alone. They suggested that other physical parameters, such as the magnetic field, may be different in the two classes. New X-ray and radio surveys, such as those by Perlman et al. (1998), Laurent-Muehleisen et al. (1998), and Caccianiga et al. (1999), have shown that the distribution of peak frequencies in the SEDs of BL Lac objects is more uniform than previously thought. What is the main physical parameter (or set of parameters) that control the SEDs of BL Lac objects?

Recent observations of the host galaxies of BL Lac objects provide new clues to understanding the physical nature of these

systems. Urry et al. (2000) and Scarpa et al. (2000) systematically studied the morphologies of BL Lac objects using the *Hubble Space Telescope* (HST) and found that there is no significant difference between the host galaxies of HBLs and LBLs. Most of the hosts are normal, giant elliptical galaxies, with no obvious evidence for ongoing mergers or strong interactions with other galaxies. Urry et al. (2000) suggest that, if the mass of the central black hole (BH) in BL Lac objects correlates with the bulge luminosity of the host galaxy, as is the case in nearby, inactive galaxies (e.g., Magorrian et al. 1998; Kormendy & Gebhardt 2001), the Eddington ratios⁵ in BL Lac objects must span a large range. A number of authors have suggested that the radio power in radio-loud quasars is controlled by the BH mass (e.g., Laor 2000; Lacy et al. 2001; McLure & Dunlop 2001; but see Ho 2002). If the host galaxies of BL Lac objects obey the BH mass/bulge luminosity relation, then the peak frequencies are evidently not controlled by the BH mass (Urry et al. 2000). This conclusion has been strengthened by the work of Falomo, Kotilainen & Treves (2002), Wu, Liu, & Zhang (2002), and Barth, Ho, & Sargent (2002a, b), who have obtained more robust BH mass estimates using the relation between BH mass and the stellar velocity dispersion of the bulge (Gebhardt et al. 2000; Ferrarese & Merritt 2000; Tremaine et al. 2002).

If BH mass is not the main parameter, what about the accretion rate? The importance of the role of accretion rates in BL Lac objects was originally recognized by Rees et al. (1982), who suggested that optically thin tori may power their central engines. A number of authors have recently revisited this issue in the context of BL Lac objects (Ghisellini & Celotti 2001;

¹ Institut für Astronomie und Astrophysik, Abt. Astronomie, Universität Tübingen, Sand 1, D-72076 Tübingen, Germany; wang@astro.uni-tuebingen.de.

² Laboratory for High-Energy Astrophysics, Institute of High-Energy Physics, CAS, Beijing 100039, P. R. China.

³ Alexander von Humboldt Fellow.

⁴ The Observatories of the Carnegie Institution of Washington, 813 Santa Barbara Street, Pasadena, CA 91101-1292.

⁵ The Eddington ratio refers to the ratio of the bolometric accretion luminosity to the Eddington luminosity, $L_{\text{edd}} \approx 1.3 \times 10^{38} m_{\text{BH}} \text{ erg s}^{-1}$, where m_{BH} is the BH mass in units of $M_{\odot}$.




FIG. 1.— Distribution of peak frequencies for our sample of BL Lac objects. Note that there is no bimodality in the distribution.

Maraschi 2001; Böttcher & Dermer 2002; Cavaliere & D’Elia 2002; Cao 2002) and more generally radio-loud quasars (Lacy et al. 2001). Using a large sample of objects which covers a wide range of nuclear activity, from nearly inactive systems to classical Seyfert 1 nuclei and quasars, Ho (2002) shows that the conventional “radio-loudness” parameter is strongly inversely correlated with the mass accretion rate.

This paper presents empirical evidence that the SEDs of BL Lac objects depend on the accretion rate.

2. THE SAMPLE

The sample of BL Lac objects is chosen according to the availability of information on (1) the luminosity of the host galaxy, (2) the luminosity of one or more emission lines, and (3) the fitted peak frequency and luminosity of the SED. The sample is limited mainly by the host galaxy observations. Relatively large ground-based optical imaging surveys of BL Lac objects have been performed by Abraham, McHardy, & Crawford (1991), Wurtz, Stocke, & Yee (1996), and Falomo & Kotilainen (1999). More recently, an *HST* survey of 132 BL Lac objects was completed by Scarpa et al. (2000) and Urry et al. (2000). Whenever possible, we have given preference to the *HST* data because of the high resolution and the uniformity of that sample. Emission-line fluxes come primarily from the studies of Stickel, Fried, & Kühn (1993), Rector et al. (2000), and Rector & Stocke (2001). Finally, we listed published values of the peak frequencies and the corresponding peak luminosities of the SEDs; these are obtained by fitting a logarithmic parabola to the multiwaveband, usually non-simultaneous, continuum.

The final sample, listed in Table 1, consists of 43 objects. For convenience, distance-dependent quantities assume $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and $q_0 = 0$. We note that the original classifications by Urry et al. (2000), which used $\log F_{\text{keV}}/F_{5\text{GHz}} = -5.5$ as the criterion to separate the two types, contained some HBLs with low peak frequencies (ν_{pk}). Figure 1 shows the distribution of ν_{pk} for our sample. There is no obvious bimodality in the distribution; some objects have “intermediate” peak frequencies. For the subsequent analysis, we somewhat arbitrarily choose $\log \nu_{\text{pk}} = 15.1$, roughly in the middle of the distribution, as the boundary between HBLs and LBLs. With this criterion, our sample has 17 HBLs and 26 LBLs.

TABLE 1
THE SAMPLE OF BL LAC OBJECTS

Source	$-M_R$	$\log \nu_{\text{pk}}$	$\log L_{\text{pk}}$	$\log L_{\text{lines}}$	Ref
0158+001	23.05	16.87	44.89	42.56	1
0205+351	23.33 ^a	17.51	44.55	42.43	1
0607+710	24.34	15.25	44.04	42.40	1
0737+744	24.32	16.16	45.21	42.65	1
0922+749	24.64	17.21	44.91	43.20	1
1207+394	24.40	16.75	45.24	42.55	1
1221+245	22.49	15.27	44.36	42.78	1
1229+643	24.07	17.23	44.60	43.39	1
1312-422	23.38 ^b	15.90 ^d	...	42.45	1
1407+595	24.78	15.90	44.50	42.77	1
1443+635	23.86 ^a	19.57	44.41	42.20	1
1458+224	23.69	15.40	45.00	44.30	1
1534+014	24.21	16.88	44.70	42.13	1
1652+398	24.20 ^c	16.00 ^e	...	42.20	2
1757+703	23.63	17.18	45.09	42.36	1
2005-489	23.86	18.26	45.47	42.03	2
2143+070	23.68	16.34	44.67	42.50	1
0122+090	23.75	13.47	44.51	42.56	1
0235+164	<27.25	13.39	46.72	44.27	2
0257+342	24.05	14.88	44.24	42.27	1
0317+183	23.71	14.36	44.27	42.60	1
0419+194	24.01	13.83	45.00	42.84	1
0537-441	<26.96	14.07	46.62	45.31	2
0851+202	<23.69	13.72	46.03	43.92	2
0954+658	<23.23	14.09	45.39	44.18	3
1144-379	<24.99	13.75	46.54	44.65	2
1235+632	23.95 ^a	14.88	44.35	42.65	1
1308+326	<27.00 ^c	13.83	46.78	44.96	2
1402+041	<23.08	14.85	44.64	43.01	1
1418+546	24.09	13.85	45.11	43.59	2
1514-241	23.54	15.09 ^f	...	42.10	2
1538+149	24.64	13.56	45.74	43.50	2
1552+202	24.39 ^a	13.70	44.74	43.51	1
1749+096	23.60	13.27	45.78	43.76	2
1749+701	<27.11	14.43	46.13	45.22	2
1803+784	<24.55	13.43	46.12	44.90	3
1807+698	23.95	14.26	44.06	43.81	4
1823+568	25.07	13.65	46.01	43.40	4
2007+777	23.89	13.66	45.48	43.76	2
2131-021	<27.02	13.33	45.67	44.17	3
2200+420	23.61	14.25	44.66	42.80	2
2240-260	<23.83	13.32	45.93	44.13	2
2254+074	24.41	13.25	44.99	43.17	2

NOTES: (a) Falomo & Kotilainen 1999; (b) Falomo & Ulrich 2000; (c) Pursimo et al. 2002; (d) Wolter et al. 1998; (e) Comastri, Molendi & Ghisellini 1995; (f) Landau et al. 1986. References for emission-line fluxes: (1) Rector et al. 2000; (2) Stickel, Fried, & Kühn 1993. (3) Rector & Stocke 2001; (4) Lawrence et al. 1996. The data for absolute magnitudes are taken from Urry et al. 2000, and the peak frequencies and luminosities are taken from Sambruna et al. 1996. The objects above and below the dividing are HBLs and LBLs, respectively.

3. STATISTICAL PROPERTIES

It is nontrivial to measure the thermal, accretion luminosity from BL Lac objects because the bulk of their emission comes from nonthermal processes associated with the jet. In this study, we adopt the strategy of using the observed luminosity of the emission lines to estimate the accretion luminosity of the disk. Emission lines themselves, however, are difficult to measure in BL Lac objects because they are both intrinsically weak and heavily diluted by the strong featureless continuum. Nevertheless, they have been detected in a number of sources, and our sample was chosen with this application in mind.

In our sample, the most commonly detected emission lines are Mg II $\lambda 2800$, [O II] $\lambda 3727$, H β , [O III] $\lambda 5007$, and H α . To estimate the *total* luminosity due to emission lines, we follow the method of Celotti, Padovani & Ghisellini (1997), who compare the luminosity of the observed lines with the fractional contribution these lines make to the total line luminosity, as




FIG. 2.— The distribution of accretion rates in BL Lac objects. The dotted histogram represent the HBLs and solid histograms show the LBLs.

determined from the relative line ratios computed by Francis et al. (1991) for a composite quasar spectrum. As in Celotti et al. (1997), we assume that the total line luminosity $\langle L_{\text{lines}} \rangle = 556 \langle L_{\text{Ly}\alpha} \rangle$. The line luminosities are given in Table 1.

With the assumption that the bulk of the line luminosity is photoionized by the accretion disk (Netzer 1990), L_{lines} should be proportional to the total disk luminosity. We define a “line accretion rate” and its dimensionless form as follows:

$$\Lambda = \frac{L_{\text{lines}}}{c^2}; \quad \lambda = \frac{L_{\text{lines}}}{L_{\text{Edd}}}. \quad (1)$$

The relation between λ and the dimensionless accretion rate $\dot{m}$ is given by equation (2). We determine BH masses from the absolute R -band magnitudes of the host galaxies, using the relation determined by McLure & Dunlop (2001) from a study of active galaxies: $\log m_{\text{BH}} = -0.5M_R - 2.96$. This relation is quite similar, but not identical, to that given by Kormendy & Gebhardt (2001) for inactive galaxies.

The above estimates of the accretion rates may carry significant uncertainties that are difficult to quantify. While the formal errors on the luminosities of individual emission lines are usually quite modest, what is less certain is the conversion to total line luminosity, as this step implicitly assumes that the relative line strengths in BL Lac objects are the same as in typical quasars. Thus, our values of L_{lines} , and hence Λ and λ , are potentially affected by systematic errors. The values of λ are additionally affected by the relatively large scatter in the BH mass versus bulge luminosity relation (~ 0.59 dex; McLure & Dunlop 2001). Despite the potentially large uncertainties on any individual measurement of Λ or λ , and their inferred accretion rates, we would like to emphasize that the *relative* values of these quantities, such as those deduced for HBLs and LBLs discussed below, should be more robust.

3.1. Accretion Rates

Figure 2 shows that the accretion rates vary over a rather wide range in LBLs, whereas they are more restricted in HBLs. The mean values of the accretion rates are $\langle \log \Lambda \rangle = 22.70 \pm 0.91$ (rms dispersion) and $\langle \log \lambda \rangle = -3.82 \pm 0.68$ in LBLs and $\langle \log \Lambda \rangle = 21.69 \pm 0.56$ and $\langle \log \lambda \rangle = -4.48 \pm 0.62$ in HBLs. These results directly show that the accretion rate may control the SEDs of BL Lac objects. It is important to note that the distributions of Λ and λ significantly overlap between HBLs and LBLs. This clearly reflects the continuous distribution of peak




FIG. 3.— The correlations among the parameters ν_{pk} , L_{pk} , and Λ . The solid and open symbols denote LBLs and HBLs, respectively.

frequencies in Figure 1.

With the assumption that the line luminosity traces the disk luminosity, the distribution of λ implies that the vast majority of BL Lac objects are highly sub-Eddington systems. In such a regime, accretion is thought to proceed via an optically thin “advection-dominated” accretion flow (ADAF; see Narayan, Mahadevan, & Quataert 1998). From Mahadevan (1997), the total luminosity from an ADAF is $L_{\text{disk}} \propto \alpha^{-2} m_{\text{BH}} \dot{m}^2$, where $\dot{m} = \dot{M}/\dot{M}_{\text{Edd}}$, $\dot{M}_{\text{Edd}} = 1.39 \times 10^{18} m_{\text{BH}}/\eta_{-1} \text{ (g s}^{-1}\text{)}$, $\eta = 0.1\eta_{-1}$ is the accretion efficiency, and α is the canonical viscosity parameter. For a reprocessing efficiency ξ , $L_{\text{lines}} = \xi L_{\text{disk}}$ and

$$\dot{m} = 2.17 \times 10^{-2} \alpha_{0.3} \xi_{-1}^{-1/2} \lambda_{-4}^{1/2}, \quad (2)$$

where $\alpha_{0.3} = \alpha/0.3$, $\xi_{-1} = \xi/0.1$ and $\lambda_{-4} = \lambda/10^{-4}$. We assume that ξ is of the same order as the covering factor of the line-emitting clouds (~ 0.1) since the total energy emitted by the clouds is simply the energy absorbed by them (Netzer 1990).

An optically thin ADAF can only exist below a critical value of the accretion rate $\dot{m}_c \approx \alpha^2$ (Narayan et al. 1998). For a plausible value of $\alpha = 0.3$, $\dot{m}_c \approx 0.1$. We find $\langle \log \dot{m} \rangle = -1.5 \pm 0.34$ for LBLs and $\langle \log \dot{m} \rangle = -2.0 \pm 0.31$ for HBLs. This indicates that BL Lac objects are in the ADAF regime, and that LBLs typically have significantly higher accretion rates than HBLs. These results are in qualitative agreement with the findings of Ghisellini & Celotti (2001) and Cavaliere & D’Elia (2002). We note that equation (2) may not be applicable to some of the LBLs with the highest values of λ because they have $\dot{m} \sim \dot{m}_c$; these objects straddle the (uncertain) transition from a standard disk to an ADAF (cf. Cao 2002).

3.2. Accretion Rates and SEDs

Figure 3 shows the dependence of the peak frequency (ν_{pk}) and peak luminosity (L_{pk}) on Λ . We find that L_{pk} correlates strongly with Λ ; the Pearson's correlation coefficient is $r = 0.76$, significant at a level of $>99.999\%$. Treating Λ as the independent variable, a linear regression fit gives

$$\log L_{\text{pk}} (\text{erg s}^{-1}) = (30.88 \pm 1.99) + (0.64 \pm 0.09) \log \Lambda. \quad (3)$$

An obvious interpretation of the $L_{\text{pk}} - \Lambda$ correlation is that the jet power is directly coupled to the mass accretion rate of the disk. The disk-jet “connection” in active nuclei has been discussed in a number of contexts (e.g., Rawlings & Saunders 1991; Falcke & Biermann 1995; Xu, Livio & Baum 1999; Ho & Peng 2001), and the present correlation provides additional support for this picture because the peak luminosity of the SED of a BL Lac object represents most of the radiated energy from the jet.

The relation between ν_{pk} and Λ is more complex. The HBL group and the LBL group each seems to define its own sequence. After excluding three strongly deviant points (0922+749, 1229+643 and 1458+224)⁶, HBLs appear to show a moderately strong correlation with $r = -0.6$ at a significance of 98%. The best-fit linear regression gives

$$\log \nu_{\text{pk}} (\text{Hz}) = (81.65 \pm 25.61) - (3.02 \pm 1.19) \log \Lambda. \quad (4)$$

The peak frequency formally scales with the accretion rate as $\nu_{\text{pk}} \propto \Lambda^{-3} \propto \dot{M}^{-6}$, but we note that this is somewhat uncertain because of the limited dynamic range in Λ . For LBLs, we have

$$\log \nu_{\text{pk}} (\text{Hz}) = (19.66 \pm 2.55) - (0.25 \pm 0.11) \log \Lambda, \quad (5)$$

with $r = -0.42$ and a significance of 97%. This implies $\nu_{\text{pk}} \propto \Lambda^{-0.25} \propto \dot{M}^{-0.5}$, much shallower than in HBLs. Bearing in mind the above caveat on the HBLs, the different slopes of the $\nu_{\text{pk}} - \Lambda$ relation may indicate differences in the physical processes associated with the jet, as discussed in the next section. It is interesting to note that HBLs and LBLs also delineate different loci, separated by a sharp break, when ν_{pk} is plotted against the radio luminosity (Costamante et al. 2001).

4. DISCUSSION

According to Xu et al. (1999), the accretion rate through the disk ($\dot{M}$) is related to the mass flux into the jet ($\dot{M}_{\text{jet}}$) by

$$\dot{M}_{\text{jet}}/\dot{M} \sim (H/R)^{2\delta-3}, \quad (6)$$

where H is the height of the disk at radius R and δ is a constant from 1.7–3.4. From the distribution of λ , we inferred that both HBLs and LBLs contain ADAFs. Since the structure of ADAFs is quasi-spherical, $H/R \sim 1$ or $\dot{M}_{\text{jet}} \sim \dot{M}$. As Blandford & Begelman (1999) argued, ADAFs inevitably lead to outflows. If energy is extracted from the spin of the BH via the Blandford-Znajek (1977) mechanism, $L_{\text{BZ}} = \epsilon_{\text{BZ}} \dot{M} c^2$, where $\epsilon_{\text{BZ}} = 1.8 \times 10^{-2}$ and the BH spin is assumed to be maximal (Armitage & Natarajan 1999). The BZ process has been strengthened by a recent observation of Fe K α line profile in MCG 6-30-15 (Wilms et al. 2001). This energy will be channeled into two parts: the kinetic luminosity $L_{\text{kin}} = \dot{M}_{\text{jet}} \Gamma c^2 / 2$ (Γ is the Lorentz factor of the jet) and the radiative luminosity L'_{rad} (seen in the comoving frame of the jet; e.g., Sikora et al. 1997). Here, we explore the possibility that the relative partition of the spin energy determines the shape of the SED. In particular, we investigate whether the different functional dependence of ν_{pk} on

Λ for HBLs and LBLs can be understood in terms of different radiative efficiencies in the jet.

Let us assume that the energy density of the random component of the magnetic field is proportional to the energy density of the relativistic electrons in the jet. Then, $B^2 \propto \langle E_e \rangle n_e$, where $\langle E_e \rangle$ is the mean energy and n_e is the number density of the relativistic electrons. With the help of $\dot{M}_{\text{jet}} \sim R_0^2 n_e m_p c$, where R_0 is the radius of the jet and m_p is the proton mass, we have $B \propto \dot{M}_{\text{jet}}^{1/2} R_0^{-1}$ and $n_e \propto \dot{M}_{\text{jet}} R_0^{-2}$. If the relativistic electrons have a power-law distribution of energies $N = N_0 E_e^{-\gamma}$, the emissivity at the peak frequency ν'_{pk} in the jet comoving frame is given by $\epsilon'_{\text{pk}} \propto N_0 B^{(\gamma+1)/2} \nu_{\text{pk}}^{(1-\gamma)/2} \propto n_e B^{(\gamma+1)/2} \nu_{\text{pk}}^{(1-\gamma)/2}$ (Pacholczyk 1970), and it follows that $\epsilon'_{\text{pk}} \propto R_0^{-(5+\gamma)/2} \dot{M}_{\text{jet}}^{(5+\gamma)/4} \nu_{\text{pk}}^{(1-\gamma)/2}$. The radiative luminosity in the jet comoving frame will then be given by $L'_{\text{rad}} \propto R_0^3 (\nu'_{\text{pk}} \epsilon'_{\text{pk}}) \propto R_0^{(1-\gamma)/2} \dot{M}_{\text{jet}}^{(5+\gamma)/4} \nu_{\text{pk}}^{(3-\gamma)/2}$. The ratio of the radiative to the kinetic luminosity will be proportional to $\Gamma^{-1} \dot{M}_{\text{jet}}^{(1+\gamma)/4} \nu_{\text{pk}}^{(3-\gamma)/2}$, which suggests that the higher peak frequencies lead to the higher radiative efficiencies.

For HBLs, we consider the limiting case that most of the energy is radiated. By setting $L'_{\text{rad}} \sim L_{\text{BZ}}$, we have $\nu'_{\text{pk}} \propto \dot{M}^{-(\gamma+1)/2(3-\gamma)}$, and the observed peak frequency is

$$\nu_{\text{pk}} = \mathcal{D} \nu'_{\text{pk}} \propto \dot{M}^{-(\gamma+1)/2(3-\gamma)} \propto \Lambda^{-(\gamma+1)/4(3-\gamma)}. \quad (7)$$

Here, we have assumed that the Doppler factor $\mathcal{D}$ does not vary strongly with $\dot{M}$. We see that ν_{pk} depends sensitively on Λ for a given γ . For a typical value of $\gamma \approx 2.6 - 2.7$ (Kirk, Melrose & Priest 1994), we get $\nu_{\text{pk}} \propto \Lambda^{-2.3}$ to $\Lambda^{-3.1}$. This is consistent with the empirical results for HBLs (eq. 4). Despite the apparent agreement, however, we must regard this result with some caution because of the extreme requirement that the bulk of the jet power emerge as radiative energy. Celotti & Fabian (1993) find that most jet sources do not radiate efficiently, although there are some objects with high radiative efficiencies (see their Fig. 2).

Since LBLs have lower peak frequencies, their radiative efficiencies are lower, and we postulate that in this case the kinetic luminosity dominates over the radiative luminosity. The maximum energy of the relativistic electrons will be given by the balance between energy loss and gain. The acceleration timescale is $\tau_{\text{acc}} \sim l(E_e)/c \propto B^{-1} E_e^n$, where $l(E_e) = l_0 E_e^n$ is the mean free path and $n = 1$ in the Bohm limit (Inoue & Takahara 1996). The timescale for energy loss due to inverse Compton scattering of external photons is $\tau_{\text{loss}} \propto (u_{\text{ext}} E_e)^{-1}$, where the energy density of external photons $u_{\text{ext}} \propto L_{\text{disk}}$ (Sikora, Begelman & Rees 1994). We thus have $E_{e,\text{max}} \propto (B/L_{\text{disk}})^{1/(n+1)}$, and the peak frequency is

$$\nu_{\text{pk}} \propto \mathcal{D} B E_{e,\text{max}}^2 \propto \dot{M}^{(n-5)/2(n+1)} \propto \Lambda^{(n-5)/4(n+1)}, \quad (8)$$

where we used $L_{\text{disk}} \propto \dot{M}^2$. For the Bohm limit ($n = 1$) or not very efficient acceleration (e.g., $n = 2$), the relation will be $\nu_{\text{pk}} \propto \Lambda^{-0.25}$ to $\Lambda^{-0.50}$. This, too, is roughly consistent with the empirical correlation for LBLs (eq. 5). In this scenario, which is similar to that discussed by Ghisellini et al. (1998) and Böttcher & Dermer (2002), the peak frequencies in LBLs are determined by cooling of relativistic electrons due to inverse Compton scattering of photons from the disk or reflected off of the broad-line emitting clouds.

⁶ The emission-line equivalent widths of 0922+749 and 1229+643 formally exceed the (admittedly arbitrary) limit of 5 Å. The value of ν_{pk} for 0922+749 was obtained from fitting only three points (Sambruna et al. 1996) and is therefore quite uncertain. There is no obvious explanation for 1458+224.

Finally, we would like to note that our analysis hinges on the assumption that the emission lines in BL Lac objects are photoionized by the disk, such that line luminosity reliably traces the accretion luminosity. Based on spectrophotometric monitoring of a sample of high-redshift, high-luminosity active galactic nuclei, Pérez, Penston & Moles (1989) find that the emission lines vary on timescales shorter than expected from photoionization models. They suggest that the broad-line region in these objects may be anisotropic. However, the variability behavior of the broad optical line $H\alpha$ in BL Lac itself appears to support that “ $H\alpha$ emission could be powered by thermal radiation from an accretion disk without significantly affecting the shape of polarization of the optical continuum” (Corbett et al. 1996, 2000). If it turns out that a significant fraction of the line luminosity is powered by the jet instead of the disk, then clearly the interpretation of Figure 3 needs to be reevaluated. However, the conclusion that BL Lac objects are highly sub-Eddington, advection-dominated systems would not change; in fact, it would only be strengthened because the inferred accretion luminosity would be even lower than we have assumed.

5. CONCLUSIONS

Using a sample of BL Lac objects with information on their host galaxies and emission-line luminosities, we show that LBLs have significantly higher accretion rates and Eddington ratios than HBLs. Both classes are highly sub-Eddington sys-

tems that may be accreting via an ADAF. We find that the peak luminosity of the SED correlates significantly with the accretion rate, lending strong support to the idea that the jet and accretion disk are closely coupled.

Furthermore, we present evidence that the peak frequency of the SED also correlates with the accretion rate, although the functional dependence between the two parameters appears to be quite different between HBLs and LBLs. We argue that these empirical trends can be qualitatively explained by invoking different radiative efficiencies in the jets of the two classes of objects. HBLs, which have lower accretion rates, evidently manage to convert more of their jet kinetic power into radiation. By contrast, in LBLs, which have higher accretion rates, a greater fraction of the jet power remains in kinetic form. However, the underlying causal connection between the radiative efficiency of the jet and the accretion rate, and the manner in which energy is channeled into relativistic electrons (Ghisellini, Celotti & Costamante 2002), remain unclear.

It would be of interest to perform spectrophotometric monitoring of nearby BL Lac objects in order to establish more conclusively the origin of their broad emission lines.

J.M.W. acknowledges support from the Alexander von Humboldt Foundation, the “Hundred Talents Program of CAS”, the Special Funds for Major State Basic Research Project and NSFC.

REFERENCES

- Abraham, R. G., McHardy, I. M., & Crawford, C. S. 1991, *MNRAS*, 252, 482
 Armitage, P. J., & Natarajan, P. 1999, *ApJ*, 523, L7
 Barth, A. J., Ho, L. C., & Sargent, W. L. W. 2002a, *ApJ*, 566, L13
 ——. 2002b, *ApJ*, submitted
 Blandford, R. D., & Begelman, M. C. 1999, *MNRAS*, 303, L1
 Blandford, R. D., & Znajek, R. L. 1977, *MNRAS*, 179, 433
 Böttcher, M., & Dermer, C. D. 2002, *ApJ*, 564, 86
 Caccianiga, A., Maccacaro, T., Wolter, A., Della Ceca, R., & Gioia, I. M. 1999, *ApJ*, 513, 51
 Cao, X. 2002, *ApJ*, 570, L13
 Cavaliere, A. & D’Elia, V. 2002, *ApJ*, 571, 226
 Celotti, A., & Fabian, A. C. 1993, *MNRAS*, 264, 228
 Celotti, A., Padovani, P., & Ghisellini, G. 1997, *MNRAS*, 286, 415
 Comastri, A., Molendi, S., & Ghisellini, G. 1995, *MNRAS*, 277, 297
 Corbett, E. A., Robinson, A., Axon, D. J., & Hough, J. H. 2000, *MNRAS*, 311, 485
 Corbett, E. A., Robinson, A., Axon, D. J., Hough, J. H., Jeffries, R. D., Thurston, M. R., & Young, S. 1996, *MNRAS*, 281, 737
 Costamante, L., et al. 2001, *A&A*, 371, 512
 Falcke, H., & Biermann, P. L. 1995, *A&A*, 293, 665
 Falomo, R., & Kotilainen, J. K., 1999, *A&A*, 352, 85
 Falomo, R., Kotilainen, J. K., & Treves, A. 2002, *ApJ*, 569, L35
 Falomo, R., & Ulrich, M.-H. 2000, *A&A*, 357, 91
 Ferrarese, L., & Merritt, D. 2000, *ApJ*, 539, L9
 Francis, P. J., Hewett, P. C., Foltz, C. B., Chaffee, F. H., Weymann, R. J., & Morris, S. L. 1991, *ApJ*, 373, 465
 Gebhardt, K., et al. 2000, *ApJ*, 539, L13
 Ghisellini, G., & Celotti, A. 2001, *A&A*, 379, L1
 Ghisellini, G., Celotti, A., Costamante, L., 2002, *A&A*, 386, 833
 Ghisellini, G., Celotti, A., Fossati, G., Maraschi, L., & Comastri, A. 1998, *MNRAS*, 301, 451
 Giommi, P., & Padovani, P. 1994, *MNRAS*, 268, L51
 Ho, L. C. 2002, *ApJ*, 564, 120
 Ho, L. C., & Peng, C. Y. 2001, *ApJ*, 555, 650
 Inoue, S., & Takahara, F. 1996, *ApJ*, 463, 555
 Jannuzi, B. T., Smith, P. S. & Elston, R. 1994, *ApJ*, 428, 130
 Kirk, J., Melrose, D. B., & Priest, E. R. 1994, in *Plasma Astrophysics*, ed. A. O. Benz, & T. J.-L. Courvoisier (Heidelberg: Springer-Verlag)
 Kormendy, J., & Gebhardt, K. 2001, in *The 20th Texas Symposium on Relativistic Astrophysics*, ed. H. Martel & J. C. Wheeler (New York: AIP), 363
 Lacy, M., Laurent-Meulessen, S. A., Ridgway, S. E., Becker, R. H., & White, R. L. 2001, *ApJ*, 551, L17
 Landau, R., et al. 1986, *ApJ*, 308, 78
 Laor, A. 2000, *ApJ*, 543, L111
 Laurent-Muehleisen, S. A., Kollgaard, R. I., Ciardullo, R., Feigelson, E. D., Brinkmann, W., & Siebert, J. 1998, *ApJS*, 118, 127
 Laurent-Muehleisen, S. A., Kollgaard, R. I., Moellenbrock, G. A., & Feigelson, E. D. 1993, *AJ*, 106, 875
 Lawrence, C. R., Zucker, J. R., Readhead, A. C. S., Unwin, S. C., Pearson, T. J., & Xu, W. 1996, *ApJS*, 107, 541
 Magorrian, J., et al. 1998, *AJ*, 115, 2285
 Mahadevan, R. 1997, *ApJ*, 477, 585
 Maraschi, L. 2001, in *The 20th Texas Symposium on Relativistic Astrophysics*, ed. H. Martel & J. C. Wheeler (New York: AIP), 409
 McLure, R. J., & Dunlop, J. S. 2001, *MNRAS*, 327, 199
 Narayan, R., Mahadevan, R., & Quataert, E. 1998, in *The Theory of Black Hole Accretion Discs*, ed. M. A. Abramowicz, G. Björnsson, & J. E. Pringle (Cambridge: Cambridge Univ. Press), 148
 Netzer, H. 1990, in *Active Galactic Nuclei*, ed. R. D. Blandford, H. Netzer, & L. Woltjer (Berlin: Springer), 57
 Pacholczyk, A. G. 1970, *Radio Astrophysics* (San Francisco: W. H. Freeman and Company)
 Pérez, E., Penston, M. V., & Moles, M., 1989, *MNRAS*, 239, 75
 Perlman, E. S., Padovani, P., Giommi, P., Sambruna, R., Laurence, R. J., Tzioumis, A., & Reynolds, J. 1998, *AJ*, 115, 1253
 Perlman, E. S. & Stocke, J. T., 1993, *ApJ*, 406, 430
 Pursimo, T., Nilsson, K., Takalo, L. O., Sillanpää, A., Heidt, J., & Pietilä, H. 2002, *A&A*, 381, 810
 Rawlings, S., & Saunders, R. 1991, *Nature*, 349, 138
 Rector, T. A. & Stocke, J. T. 2001, *AJ*, 122, 565
 Rector, T. A., Stocke, J. T., Perlman, E. S., Morris, S. L., & Gioia, I. M. 2000, *AJ*, 120, 1626
 Rees, M. J., Begelman, M. C., Blandford, R. D., & Phinney, E. S. 1982, *Nature*, 295, 17
 Sambruna, R. M., Maraschi, L., & Urry, C. M. 1996, *ApJ*, 463, 444
 Scarpa, R., Falomo, R., & Pian, E., & Treves, A. 1995, *A&A*, 303, 730
 Scarpa, R., Urry, C. M., Falomo, R., Pesce, J. E., & Treves, A. 2000, *ApJ*, 532, 740
 Sikora, M., Begelman, M. C. & Rees, M. J. 1994, *ApJ*, 421, 153
 Sikora, M., Madejski, G., Moderski, R., & Poutanen, J. 1997, *ApJ*, 484, 108
 Smith, P. S. & Sitko, M. L., 1991, *ApJ*, 383, 580
 Stickel, M., Fried, J. W., & Kühr, H. 1993, *A&AS*, 98, 393
 Tremaine, S., et al. 2002, *ApJ*, in press (astro-ph/0203468)
 Urry, C. M., & Padovani, P. 1995, *PASP*, 107, 803
 Urry, C. M., Scarpa, R., O’Dowd, M., Falomo, R., Pesce, J. E., & Treves, A. 2000, *ApJ*, 532, 816
 Wilms, J., et al., 2001, *MNRAS*, 328, L7

Wolter, A., et al. 1998, A&A, 335, 899

Wu, X.-B., Liu, F. K., & Zhang, T. Z. 2002, A&A, 389, 742

Wurtz, R., Stocke, J. T., & Yee, H. K. C. 1996, ApJS, 103, 109

Xu, C., Livio, M., & Baum, S. A. 1999, AJ, 118, 1169